

# What You Need To Know About Your Medicine For Tuberculosis (TB):

## Isoniazid (INH) And Rifapentine (RPT)

### About Your Medicine:

ISONIAZID (eye-soe-NYE-a-zid) and RIFAPENTINE (rif-a-pen-tine) are antibiotics used in combination to treat tuberculosis (TB) infection.

### Before Using This Medicine:

Tell your doctor, nurse, and pharmacist if you:

- Have drug allergies
- Are pregnant or plan to become pregnant
- Are breastfeeding
- Are taking any other prescription medicines
- Are taking over-the-counter medicines (such as cold medicines, pain relievers, vitamins, herbs, etc.)
- Have any other medical problems including diabetes, gout, liver disease, or kidney disease

### Tips to Help You Take Your Medicine:

- Take exactly as directed.
- Do not miss any doses or take two doses at the same time.
- Do not share your medicines with others.
- Keep this and all other medicines out of the reach of children.
- Store in a cool, dry, place away from extreme heat or cold.

[www.scdhec.gov/tb](http://www.scdhec.gov/tb)



South Carolina Department of Health  
and Environmental Control

### Warnings:

- DO NOT drink alcohol while on this medicine to avoid possible liver problems.
- If you think you have taken an overdose of this medicine, call Poison Control at 1-800-222-1222.

### Possible Side Effects of TB Medicines:

- Upset stomach or stomach pain
- Nausea or vomiting
- Severe weakness or tiredness
- Flu-like symptoms
- Coffee/tea colored urine or light stools
- Yellow eyes or skin
- Rash or itching
- Easily bleeding or bruising
- Blurred vision
- Pain or tingling in hands, feet, or joints
- Dizziness
- Fever

### INH may occasionally cause:

- Dizziness
- A tingling sensation or numbness of mouth, hands, and feet

### RPT may:

- Interfere with birth control pills/implants and methadone treatment
- Cause urine, sweat, or tears to turn orange
- Permanently stain soft contact lenses
- Cause sensitivity to the sun



Please contact  
your DHEC nurse  
immediately if you  
experience any of  
these side effects.

# သင့်၏ (TB) အဆုတ်ရောဂါဖို့ ဆေးက သိထားရန်ဘာလို့လဲ။

## Isoniazid (INH) And Rifapentine (RPT)

### သင့်ဆေးအဖော်ကား။

ISONIAZID (eye-soe-NYE-a-zid) နှင့် RIFAPENTINE (rif-a-pen-tine) သည် အဆုတ်ရောဂါ (TB) ကူးစက်ပြီး အဆုတ်ရောဂါ (TB) စင်္ကြာပုန့်ပေါင်းကုသရန်အတွက် ပြုစုပါသည်။

### ဤဆေး အသုံးပြုပုံစံ။

သင့်၏ ဆရာဝန်၊ သူနာပြုဆရာမ နှင့် ဆေးပညာရှင်များက ပျော်ပျော်ပါးပါး အကယ့်သင့်မှာ။

- ဆေးပွင့် မတည့်စေပါ။
- ကျွမ်းဝင်မှုရှိ(သို့) ကျွမ်းဝင်မှုမရှိမိန်းမကလေးများ။
- မိခင်က ကလေးကို ဂရုစိုက်ပါ။
- ဆရာဝန်ပေးသော ဆေးလက်မှတ်နှင့် ဆေးသောက်ပါ။
- ခုန့်မှတ်သော ဆေးပုံစံသည် ( ခေါ်ကုသဆေး၊ နာကျင်သကဲ့သို့သော ဆေး၊ အားဆေး၊ ဆေးဖက်ဝင်အပင်များ တို့ဖြစ်ပါသည်။ )
- တျားသော ကန့်သတ်ချက်များရှိသော ဆေးဝါးများ၊ အဆုတ်ရောဂါ၊ အသက်ရှူရောဂါ၊ ကော်ကုန်ပုရောဂါ တို့ဖြစ်ပါသည်။

### သင့်ဆေးသောက်ရန် အဆင်ပြေအောင်။

- အညှင်းအထိုး သောက်ပါ။
- ကော်သက်ခြင်း (သို့) ပုံစံခါစာကုတ်စုခါတည်းသောက်ခြင်းမျိုးပါ။
- သင့်ဆေးကိုတျားသုံးနေသော ဝေဒနာကျေအောင် ပျော်ပါ။
- ဤဆေးပွင့် တျားသော ဆေးများကိုလေးမယူမီသောနေ့က သိမ်းထားပါ။
- အေးချမ်း ပျော်ရွှင်သော ပုံစံမျိုး မပျက်စီးအောင်လည်းကောင်း၊ နေရာကို သိမ်းထားပါ။

[www.scdhec.gov/tb](http://www.scdhec.gov/tb)



### သတိပေးချက်များ။

- ဤဆေးသောက်နေခိုက်တွင် အသက်ပျောက်မှုများ ပေါ်ပေါက်ပါသည်။ အရက်သောက်စားခြင်း မပြုပါ။
- အကယ့်ဤဆေးကိုအသုံးပြုသည့် သင့်ထံသို့ အဆီပျော်စပ်သုံးသုံးရန်နှင့် ဤဖုန်းနံပါတ် 1-800-222-1222 ဆက်ပါ။

### ဆက်သွယ်ရန်သော (TB) ဆေးများ။

- ဝမ်းဗိုက်အပင် (သို့) ဝမ်းဗိုက်နာကျင်မှု။
- မူးယစ်စွမ်း (သို့) ပိုမိုအနံ့ရှိသည်။
- အားနည်းခြင်း (သို့) ပင်ပန်းခြင်း ဖြစ်နိုင်ပါသည်။
- တုတ်ကေးပြင်သို့လက်ကောက်ရခြင်း။
- မစင်သောဆေးရည် ကော့ဖီနှင့် လက်ကောက်ရည်အရောအနှော။
- မိကျွန်း အရေပြားက အဝါရောင်ရှိသည်။
- ရမ်းပျောင်း (သို့) အဖုတို့ဖြစ်ပေါ်နိုင်ပါသည်။
- အလျှော့တက် သေခြင်းဖြစ်နိုင်ပါသည်။ (သို့) ညီမည်းရောင်ရှိပါသည်။
- မိကျွန်း မီးခိုးရှိပါသည်။
- ကျွမ်းဝင်မှုရှိ (သို့) အဆုတ်ရောဂါ နာကျင်မှု။
- မူးယစ်ခြင်းဖြစ်ပါသည်။
- ကျွမ်းဝင်မှုရှိပါသည်။

### INH ပုံစံသော အဖော်ကားများ။

- မူးယစ်မှုပုံစံများ။
- ခံနိုင်ရည်မရှိမှုပုံစံများ (သို့) ပါစပေါ့၊ ပျော်၊ လက် ကိုင်ရန်ဖြစ်ပါသည်။

### RPT ပုံစံသော က။

- သင့်တော်သောဆေးပုံစံ၊ သင့်တော်သောအပွင့် ကုသဆေးပုံစံဖြစ်ပါသည်။
- ဆီးရေ၊ ခေါ် (သို့) မိကျွန်းများ အဝါရောင်ထွက်နိုင်ပါသည်။
- အခွံရောင် ပေါ်ပေါက်သောအရာနှင့် မွန်ဘီလူးကုန်ပါ။
- နေရောင်ခြည်ကိုလက်ပြုလုပ်ခြင်း ခံနိုင်ပါသည်။



အကယ့်ဤ နှစ် ကိုခံစားမိသော သင့်၏ စားရပါက သင့်၏ DHEC ဆရာမကို ခံကုန်ဆက်သွယ်ပါ။